

**POSTERS****Table of Contents**

67	Molecular mechanism of inflammation-related diseases	208	Synthetic biopolymers for biomedicine
92	DNA variation	215	Integrative approaches to structural and synthetic biology
103	Cardiovascular diseases	219	Induced pluripotent cells
114	Intracellular ion channels and transporters	221	Long noncoding RNA
121	RNA processing	222	Neurodegeneration
123	Signal transduction	234	Cell therapy and regenerative medicine
135	Mitochondria and signaling	243	Small noncoding RNA
147	DNA architecture	248	Proteins: structure, disorder and dynamics
153	RNA transcription	279	Plant biotechnology
156	DNA editing and modification	289	Natural networks and systems
160	RNA transport and translation	292	RNA in pathogenesis and therapy
165	Single cell analysis and imaging	294	Molecular biology of aging
171	Calcium and ROS signalling	301	Plant–environment interaction
175	Sulfur metabolism and cellular regulation	312	Synthetic networks and systems
179	Molecular neurobiology	313	Multicomponent complexes
188	RNA turnover	318	Cell signaling in tumor biology
191	Cytoskeleton and molecular mechanisms of motility	371	Bionanotechnology
198	Rare diseases	391	Epigenetics and protein glycosylation
204	Signaling in brain cancer	396	Genome editing (CRISPR)
		402	Proteomic technologies
		408	Education, training, and career planning in molecular life sciences
		409	General topics – various

Abstracts submitted to the 44th FEBS Congress, taking place in Krakow, Poland from 6th to 11th July 2019, and accepted by the Congress Organizing Committee are published in this Supplement of *FEBS Open Bio*. Late-breaking abstracts are not included in this issue.

**About these abstracts**

Abstracts submitted to the Congress are **not peer-reviewed**. In addition, abstracts are published as submitted and are **not copyedited** prior to publication.

We are unable to make **corrections of any kind** to the abstracts once they are published.

**Indexing**

Abstracts published in *FEBS Open Bio* Supplement for the 44th FEBS Congress will be included individually in the Conference Proceedings Citation Index published by Web of Science.

**How to cite these abstracts**

AuthorOne, A., AuthorTwo, B. (2019). Abstract title. *FEBS Open Bio*, 9: Abstract number\*. doi:10.1002/2211-5463.12675

\* Each poster has been given a unique number beginning with the letter P; the next part relates to the session in which the poster will be presented.

P.01	Molecular mechanism of inflammation-related diseases	P.22	Induced pluripotent cells
P.02	DNA variation	P.23	Long noncoding RNA
P.03	Cardiovascular diseases	P.24	Neurodegeneration
P.04	Intracellular ion channels and transporters	P.25	Cell therapy and regenerative medicine
P.05	RNA processing	P.26	Small noncoding RNA
P.06	Signal transduction	P.27	Proteins: structure, disorder and dynamics
P.07	Mitochondria and signaling	P.28	Plant biotechnology
P.08	DNA architecture	P.29	Natural networks and systems
P.09	RNA transcription	P.30	RNA in pathogenesis and therapy
P.10	DNA editing and modification	P.31	Molecular biology of aging
P.11	RNA transport and translation	P.32	Plant–environment interaction
P.12	Single cell analysis and imaging	P.33	Synthetic networks and systems
P.13	Calcium and ROS signalling	P.34	Multicomponent complexes
P.14	Sulfur metabolism and cellular regulation	P.35	Cell signaling in tumor biology
P.15	Molecular neurobiology	P.36	Bionanotechnology
P.16	RNA turnover	P.37	Epigenetics and protein glycosylation
P.17	Cytoskeleton and molecular mechanisms of motility	P.38	Genome editing (CRISPR)
P.18	Rare diseases	P.39	Proteomic technologies
P.19	Signaling in brain cancer	P.40	Education, training, and career planning in molecular life sciences
P.20	Synthetic biopolymers for biomedicine	P.41	General topics – various
P.21	Integrative approaches to structural and synthetic biology		

differentiation, cell proliferation, embryonic development. Even slight shift in miRNA level could lead to significant changes of transcriptome, and in a result of cell phenotype. In the last decade, over 2500 human mature miRNA sequences were deposited in miRBase. The function of many of them have been found and anti-miRNA, as potential therapy tools have been designed. Despite of an enormous data of miRNA, there are still many questions concerning miRNA function to be solved. The generally accepted model of the miRNA-guided RNA down-regulation suggests that mature miRNA targets mRNA in a nucleotide sequence-specific manner. Recently, we analyzed the nucleotide content of human mature miRNAs and showed that the most abundant nucleotide in miRNAs is guanosine. We identified guanosine-rich miRNAs and found that the level of G in some miRNAs is even over 90%. We noticed that some miRNAs (179 of 2565 human ones) have quadruplex specific motif GGN(1-7)GGN(1-7)GGN(1-7)GG. Furthermore, we found that at least 50% of miRNAs may adopt tertiary structure. Using specific nucleases, NMR, UV/Vis and CD spectroscopies, small-angle X-ray solution scattering (SAXS) as well as and molecular dynamics, as the first ones, we gave evidences that some miRNAs may adopt quadruplexes, e.g. miR-3620, 4507, hairpin and/or homoduplex structures, e.g. miR-21, miR-93 and miR-296. We also showed that the structure of miRNA has functional consequences and suggested that miRNAs may function also beyond RISC and are even more sophisticated regulators, that it was previously expected. We think that the knowledge of the miRNA structure and its dynamic may give a new insight into miRNA-dependent gene regulation mechanism and be a step forward in the understanding their function and involvement in cancerogenesis.

#### P-26-003

##### Bacterial 6S RNAs under stress conditions

D. Elkina<sup>1</sup>, O. Burenina<sup>2</sup>, V. Bannikova<sup>2</sup>, E. Kubareva<sup>1</sup>

<sup>1</sup>A.N. Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University, Moscow, Russia,

<sup>2</sup>Department of Chemistry, Lomonosov Moscow State University, Moscow, Russia

6S RNA is a highly abundant small non-coding RNA found in diverse classes of prokaryotes, including bacteria with an extremely small genome, that indicates an important role of 6S RNA in the cell. 6S RNA has a unique secondary structure which mimics DNA promoter, thus 6S RNA is able to bind RNA polymerase (RNAP) holoenzyme and inhibit transcription in case of an unfavorable environment. Moreover, DNA-dependent RNAP can use 6S RNA as a template for synthesis of short pRNAs, when it needs to release from 6S RNA and proceed transcription of genes. Under normal conditions 6S RNA knockout does not affect *E. coli* growth, while it becomes essential for cell adaptivity to starvation and alkaline stress. Some other bacteria express 6S RNA at highest levels under standard cultivation conditions. These facts raise a number of questions, like (i) what is the mechanism of 6S RNA-dependent transcription inhibition (and its retraction) in these bacteria, (ii) whether 6S RNA is involved in some stress responses, and (3) what type of genes are affected. In the present work we characterized properties and functions of 6S RNAs from *B. subtilis* and *R. sphaeroides* in comparison to *E. coli*. It was shown that in *B. subtilis* 6S RNA knockouts leads to widespread changes in cellular transcriptome and proteome and results into high cell viability under alkaline conditions. For the first time, *R. sphaeroides* 6S RNA was characterized and osmotic and oxidative phenotypes of cells with deletion of 6S RNA gene were found. Surprisingly, decreased viability of *E. coli* 6S RNA knockout strain was observed in case of response to one of the stress, and some genes affected by the lack of 6S

RNA under these conditions were revealed by qRT-PCR. Thus, for the first time we experimentally proved a great variety of 6S RNA functions in different bacteria and its involvement into diverse cellular stress responses. This work was supported by the Russian Foundation for Basic Research (project No. 19-04-00791).

#### P-26-004

##### MIR3648 and MIR3867 target heterochromatin

A. Gavrilov<sup>1</sup>, A. Zharikova<sup>2</sup>, A. Galitsyna<sup>1,2,3,4</sup>, A. Mironov<sup>2,4</sup>, S. Razin<sup>1</sup>

<sup>1</sup>Institute of Gene Biology, Russian Academy of Sciences, Moscow, Russia, <sup>2</sup>Faculty of Bioengineering and Bioinformatics, Lomonosov Moscow State University, Moscow, Russia, <sup>3</sup>Skolkovo Institute of Science and Technology, Moscow, Russia, <sup>4</sup>Institute for Information Transmission Problems, Russian Academy of Sciences, Moscow, Russia

Non coding RNAs (ncRNAs) participate in various biological processes from regulating enzymatic activities to sustaining spatial organization of the genome. MicroRNAs, whose canonical function is suppressing mRNA translation in the cytoplasm via RNA interference, occur in the nucleus as well where these miRNAs may pair with other ncRNAs localized in certain genome regions and trigger repression or activation of these regions. Here we studied RNA-DNA interactome of the human erythroleukemia cell line K562 by a new method, which is based on adapter-mediated RNA-DNA ligation in cross-linked nuclei followed by sequencing. We identified several hundreds of ncRNAs enriched in active or repressed chromatin. Of particular interest are MIR3648 and MIR3867. We found that these miRNAs establish contacts genome-wide and are ranked among the first in localization to repressed chromatin and the inactive spatial chromatin compartment among all ncRNAs identified in the study. MIR3648 and MIR3867 favor bulk heterochromatin over Polycomb repressed chromatin. They associate with regions of late replication and lamina associated domains, are depleted from the bodies of transcribed genes and enriched in gene deserts. The frequency of contacts of MIR3648 and MIR3867 with gene poor chromosome 18 is about 2 times higher than genome average. Interestingly, MIR3648 and MIR3867 genes are hosted within the 5' external transcribed spacer of the 45S rRNA operon. Importantly, the rRNA itself does not show preferences for active or repressed chromatin. Based on the above observations we speculate that MIR3648 and MIR3867 play a role in heterochromatin formation at the genome scale. Of note, previous studies have revealed multiple examples of miRNA involvement in transcriptional silencing of individual genes. However, MIR3648 and MIR3867 represent the first example of miRNAs associated with inactive chromatin genome-wide. This work was supported by the Russian Science Foundation (grant 18-14-00011).

#### P-26-005

##### Inhibition of SIRT1/p53 pathway by miR-211 and induction of cell death in breast cancer cells

M. Nourbakhsh\*, S. Yarahmadi\*, Z. Abdolvahabi, Z. Hesari,

M. Tavakoli-Yaraki, Z. Yousefi, P. Seiri

*Iran University of Medical Sciences, Tehran, Iran*

Sirtuin1 (SIRT1), a class III histone deacetylase enzyme, can deacetylate various proteins including p53 and regulate diverse physiological and pathological processes. SIRT1 has been suggested as a promoting factor in tumor development and