

Earth and Planetary Science Letters 179 (2000) 237-242

EPSL

www.elsevier.com/locate/epsl

First direct dating of Late Pleistocene ice-wedges by AMS

Yurij K. Vasil'chuk^{a,b,*}, Johannes van der Plicht^c, Hogne Jungner^d, Eloni Sonninen^d, Alla C. Vasil'chuk^{a,b}

^a Cryolithology Department, Faculty of Geography, Lomonosov's Moscow University, Vorob'yovy Hills, 118899 Moscow, Russia

^b Theoretical Problems Department, The Russian Academy of Sciences, Denezhnyi per. 12, 121002 Moscow, Russia

^c Radiocarbon Laboratory, Centre for Isotope Research, Groningen University, Nijenborgh 4, 9747 AG Groningen, The Netherlands ^d Dating Laboratory, Helsinki University, P.O. Box 11, FIN-00014 Helsinki, Finland

Received 27 April 1999; received in revised form 10 April 2000; accepted 11 April 2000

Abstract

We present the first direct dating by ¹⁴C-accelerator mass spectrometry of three Late Pleistocene syngenetic ice-wedges from the Seyaha cross-section. They are representative of permafrost with multistage ice-wedges from the North of Western Siberia. The most important result is the clear vertical age stratification of the ice, i.e. the old ice is located beneath the young. This shows that a timescale can be assigned to these ice-wedges penetrating down into the permafrost. The age of the ice shows a depth of not more than 3-5 m for frost cracking; water penetrated into the ice-wedges at that depth. The lower part of the ice-wedges from the Seyaha cross-section has been dated between 21 000 and 14 000 BP. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: accelerator mass spectrometry dating; ice wedges; Pleistocene; paleoclimatology

1. Introduction

Syngenetic ice-wedges, a type of ground glaciation, occur throughout areas covered with permafrost. Ice from such ice-wedges forms a unique archive, preserving many features of the past environment. The analysis of the constituents collected in the ice provides new information about the history of the climatic system.

Syngenetic ice-wedges as old as 40 000–50 000 years are available for investigation in natural exposures along sea shores, river or lake banks and

* Corresponding author. Tel.: +7-095-939-3673;

in quarries [1,2]. This allows detailed and clean sampling of the ice, independent of its age. Syngenetic ice-wedges can be dated either by conventional dating of bulk organic material from the host sediments surrounding the ice, or (almost) directly by accelerator mass spectrometry (AMS) dating of organic micro-remains, which were transported by aerosols and trapped in the icewedge ice. Earlier attempts to date macro-organic remains directly from ice-wedges yielded conflicting results [2].

The peculiarities of ice-wedge dating result from their formation process [1]. The ice-wedges result from the filling of frost cracks with snowmelt water, consisting of many vertical veins. Syngenetic ice-wedges form under conditions of slow, continuous sedimentation accompanied by re-

Fax: +7-095-114-4518; E-mail: vasilch@orc.ru

⁰⁰¹²⁻⁸²¹X/00/\$ – see front matter © 2000 Elsevier Science B.V. All rights reserved. PII: S 0 0 1 2 - 8 2 1 X (0 0) 0 0 1 2 2 - 9

peated frost cracking. Subaqueous and subaerial stages of the wedges can be distinguished by their sedimentation conditions. The subaqueous stage corresponds to a relatively short period (about 1000 years) of a high accumulation rate and the subaerial stage corresponds to a relatively long period (more than 5000 years) of a low accumulation rate. The growth of ice-wedges is a fluctuating process, which is slower during subaqueous deposition of sand, sandy loam or loam and faster in subaerial conditions during accumulation of peaty sediments or peat. The ¹⁴C dates of the host sediments are almost all from such subaerial stage.

Direct dating of ice-wedges gives information about the exact depth of frost cracking. This can be done by dating small amounts of organic material by means of AMS [1,3]. Here we report on such a study for a representative ice-wedge cross-section: Seyaha in the North of Western Siberia.

2. The Seyaha cross-section

The site at the Seyaha Settlement on the east coast of Yamal Peninsula (70°N, 72°E) was investigated by the authors in 1978, 1980 and 1996. The exposure extends more than 4 km along the coastline of the Ob Bay. This sequence with a depth of 22–24 m is especially valuable for paleo-geographical reconstruction thanks to the abundance of multistage ice-wedges. The sediments accumulated continuously during almost 20 000 years: from 30 000 to 11 000 BP (according to ¹⁴C dates of the host sediments). Three characteristic parts can be observed in the geocryologic structure of the section (Fig. 1).

The lowest 11 m is characterized by 3 m wide ice-wedges of pure ice. This part of the sediment formed when the lower coastal parts of the Ob Bay were inundated, as indicated by the structure of these layers, with organic inclusions. The host sediments accumulated in alternating subaqueous-subaerial conditions, they are laminated and contain organic material such as rootlets, stems and leaves (subaerial layers) with sandy Depth



Fig. 1. Syngenetic sediments with ice-wedge complexes near the Seyaha settlement. The AMS dates are indicated. Sample No. 1 is for re-deposited material (see text). Legend: (1) sand; (2) loamy sand with small content of allochthonous organic material; (3) horizons enriched with autochthonous and allochthonous organic material; (4) syngenetic Late Pleistocene large ice-wedges; (5) AMS-samples of non re-deposited micro-organic remains from syngenetic Late Pleistocene ice-wedges (numbers correspond with Table 1) and (6) marine foraminifera.

loam admixture (subaqueous layers). The main part of the organic material consists of moss remains of Drepanocladus fluitans (Hedw.) Warnst, Tomentipnum nitens, Paludella squarrosa, Scorpidium scorpioides, herbs Eriphorum vaginatum, Carex caespitosa, Carex rostrata, Carex inflata and also single remains of shrubs Oxycoccus sp. and Ledum palustre L. In addition, organic laminae contain a small amount of mineral admixture.

The middle 8-9 m of the section contains 1-1.5 m wide ice-wedges. These wedges also penetrate into the ice-wedge body of the lower stage and show vertical layers containing mineral inclusions. The host sediments are laminated but have a low organic content, because we believe that this part of the sediment accumulated during a period of higher water levels of the Ob Bay.

The uppermost 2–3 m consists of laminated yellow sand (subaqueous layers). This part of the sediment is assumed to have formed during a more inundated period. This conclusion is shown by both structural features of the deposit and from the presence of well-preserved and not

re-deposited sub-saline foraminifera: *Elphidium* subclavatum Gudina, Pninaella pulchela Parker, Protelphidium parvum Gudina, Discorbis sp., Miliolinella cf. subrotunda (Montagu), Discorbis deplanatus Gudina, Globulina glacialis Gushman et Ozawa, etc. The uppermost part of the section contains 1.5 m wide ice-wedges; these penetrate into the ice-wedge body of the middle stage. These ice-wedges also contain vertical inclusions of sand.

Previous work on the chemical composition of ice from ice-wedges and segregated ice (ice lenses in the host sediments) shows a varying influence of the Ob Bay for different sedimentation regimes. For a more detailed description we refer to [1].

3. Results and discussion

The age of Late Pleistocene syngenetic icewedges in the Seyaha cross-section has been determined in two ways, by conventional radiocarbon dating and by AMS dating.

3.1. Conventional dating of host sediments

Radiocarbon dating of organic material from the host sediments was done in two stages. At first, a series of six radiocarbon dates ranging

from 30000 to 22000 BP from the host sediment were obtained from the Institute of Geology, Moscow (laboratory code GIN) in 1982 [4]. The lower part of the section was dated, which consisted of laminated organic material with sandy loam. Later, the lower part of the section was re-dated and also several very important dates from the host sediment of the upper part were obtained in 1996-1997 from the University of Helsinki (laboratory code Hel). There is a good agreement between radiocarbon dates from the same horizons, which have been obtained in 1980 and in 1997 (after the field seasons 1979 1996). For example, the date and of $30\,100\pm1500$ BP (GIN-2477) was obtained at 21 m depth at the basement of the ice-wedge complex and the date of 29500 ± 400 BP (GIN-8936) was obtained 17 years later at a depth of 20 m (Table 1). The date of 22600 ± 600 BP (GIN-2475), from the depth of 10 m, is very close to the date of 22510 ± 330 BP (GIN-8931), obtained 17 years later from a depth of 11 m [3]. The dates are very consistent in spite of the possible presence of allochthonous components in the sediments. In particular, the bottom and top of the lowermost 11 m of the laminated sediment are well dated. The bottom section was first dated to about 30 000 BP. One AMS date of a Betula nana twig from the same sample gave an age of

Table 1

¹⁴C dates for the Seyaha ice-wedge complex, Russia: dating of the host sediment

| Sample | Depth (m) | Material | Laboratory code | ¹⁴ C age | |
|---------|--------------|----------|-----------------|---------------------|--|
| | | | | (yr BP) | |
| 363/55 | 0.8 | peat | Hel-3942 | 11620 ± 90 | |
| 363/77 | 3.2 | peat | Hel-4023 | 17290 ± 250 | |
| 279/10 | 8.6 | peat | GIN-2473 | 22700 ± 300 | |
| 279/13 | 10.0 | peat | GIN-2475 | 22600 ± 600 | |
| 363/207 | 11.0 | peat | GIN-8931 | 22510 ± 330 | |
| 363/111 | 12.0 | peat | Hel-4046 | 22850 ± 440 | |
| 279/16 | 12.0 | peat | GIN-2474 | 23500 ± 400 | |
| 363/206 | 12.0 | peat | Hel-4043 | 24460 ± 650 | |
| 363/62 | 12.2 | peat | Hel-4056 | 25300+900 | |
| 363/112 | 15.2 | peat | Hel-3943 | 27890 ± 90 | |
| 279/23 | 16.2 | peat | GIN-2476 | 24300 ± 300 | |
| 363/212 | 20.5 | peat | GIN-8936 | 29500 ± 400 | |
| 279/28 | 20.9 | peat | GIN-2477 | 30100 ± 1500 | |
| 363/211 | 20.9 | twig | Hela-201 | 31200 ± 900 | |
| 363/208 | 20.9 | peat | Hel-3950 | 36800+3300/-2100 | |

31 000 BP (Hela-201). This confirms that the lowermost 11 m layer began to form about 30 000 BP. The dates for samples taken from different sites of the 4 km long exposure along the Ob Bay coastline from the same depth suggest the chronology. Five dates from 11 m depth were obtained, ranging from 22 300 to 22 800 years BP.

The radiocarbon dates of the peat (Table 1) show that the 11 m of sediment at the bottom of the cross section accumulated during ca. 7500 years. The accumulation rate was about 1.3 m per 1000 years.

The accumulation rate and age of the sediments can be used to obtain a timescale for the ¹⁸O curves of the ice-wedge ice [1,4,5]. However, a more accurate age can be obtained by direct AMS radiocarbon dating of organic material extracted from ice-wedge ice. When only dates of organic material from host sediments surrounding ice-wedges were available, we had assumed that ice-wedges at the bottom of the cross-section began to grow about 27 000 years ago [1,4]. We conclude that we have underestimated the possibility of re-deposition of old organic material under syngenetic permafrost conditions.

3.2. Direct AMS dating of ice-wedge ice

In 1996, large samples of ice from three different levels of ice-wedges were collected. Extracted organic material was radiocarbon-dated using AMS. Up to 10 kg of ice was sampled and thawed in a closed bottle near the sampling site. After settling during 48 h, the water was discharged and the residue collected.

The ice-wedges themselves were dated by small organic remains obtained from the ice. The mate-

rial was pre-treated by the usual acid–alkaline– acid method [6], combusted to CO_2 and subsequently reduced to graphite and measured by AMS [7] in Groningen (laboratory code GrA).

The AMS dates show that ice-wedges began to form about 21 000 BP (sample No. 3 in Fig. 1). From these direct dates of ice-wedge ice from 21 000 to 14 000 BP we are able to deduce that the 1980 host sediment samples No. 10, 13 and 16 (GIN-2473, 2474, 2475) must have been contaminated by older organic material. Sample No. 23 (GIN-2476) from the same collection contained autochthonous organic material, yielding a more consistent result. It is likely that contemporaneous ice-wedge ice is found 3–5 m lower, because water penetrated into ice-wedges at that depth (the depth of frost cracking at that time).

From the AMS dates for the deepest samples (see Fig. 1, No. 2 and 3), we conclude that the icewedge vertical accumulation rate was about 1.2-1.3 m per 1000 years. The soluble alkaline extracts are older than the organic component for samples 1 and 2 (Table 2). These differences can be explained by the formation of ice-wedges of the higher stages 2 and 3 in open beach conditions. The ice-wedges could have been contaminated with fine organic dust, which blew into frost cracks from neighboring exposures of ancient sediments. Some organic material, which penetrated into the frost cracks may have come from the sand with a high concentration of ancient organic material. Evidence for this suggestion is the high concentration of re-deposited pollen and spores older than Pleistocene in the upper part of the section. However, the AMS dates of both the alkaline extract and the organic material of the lowest sample (No. 3) are almost identical.

Table 2

¹⁴C dates for the Seyaha ice-wedge complex, Russia: direct dating of the ice-wedge

| e dates for the segund fee wedge complex, Russia, direct dating of the fee wedge | | | | | | | |
|--|---------|--------------|-----------------------------------|------------------------|------------------------------------|--|--|
| No. | Sample | Depth (m) | Material | Laboratory code | ¹⁴ C age (yr BP) | | |
| (1) (1a) | 363/27 | 1.8 | micro-organic alkaline extract | GrA-10538 GrA-9847 | 14550 ± 100 19920 ± 130 | | |
| (2) (2a) | 363/87 | 12.0 | micro-organic alkaline extract | GrA-10539 GrA-9848 | 14720 ± 100 23620 ± 160 | | |
| (3) (3a) | 363/125 | 20.6 | micro-organic alkaline extract | GrA-10536 GrA-10535 | 20960 ± 140 21440 ± 140 | | |
| | | | | | | | |

The natural conditions during ice-wedge formation at the lowest stage were less affected by contamination because during that time there was tundra vegetation and peat covering the seacoast. This young organic material penetrated into the frost cracks at these times, which explains why the dates of both fractions for sample No. 3 are identical.

Up to now, ice-wedges have been dated by ¹⁴C dating the surrounding host sediments. The correlation between these dates and ¹⁸O data of the ice, yielding paleoclimate information, is difficult, for example, if the depth of frost cracks is different for the different epochs.

Whether the 14 C dates of micro-organic material included in the ice-wedge are correct, depends on whether they have an autochthonous or allochthonous origin [9]. The dates of alkali-soluble fractions of the 14 C sample are older, apparently because old organic material is re-deposited into the initial cracks. For the Seyaha cross section presented in this study, it is possible to select samples with a high content of autochthonous material.

The AMS dates are crucial for the investigation of the changes of levels in the Ob Bay during the late Pleistocene. The upper half of the section accumulated between ca. 11000 and 19000 BP. The conventional date at 17000 BP represents the time of the deepest regression [8]. At 11000 BP the subaqueous stage terminated. It may be possible to verify our dating of ice in the icewedges by AMS dating of CO_2 trapped in the ice [10].

4. Conclusions

For the first time, we have directly dated icewedges by ¹⁴C-AMS, using micro-remains of organic material extracted from the ice. Syngenetic ice-wedges formed by the penetration of melt water in frost cracks in the ice-wedges, contemporaneously with sediment accumulation. The older ice is located below the younger, confirming stratigraphy. Our data show a good perspective for dating Late Pleistocene syngenetic ice-wedges by means of radiocarbon measurements. The time of formation of ice-wedges, determined by direct (AMS) and indirect ice dating (host sediments which can be dated by conventional 14 C) is consistent. We can apply these methods to other important cross sections in the permafrost. In combination with stable isotope (18 O, 2 H) analysis of the ice it now becomes feasible to reconstruct chronologies for winter paleotemperatures of ground ice.

Acknowledgements

We would like to thank Nadine Budantseva, Leopold Sulerzhitsky and Julia Chizhova for invaluable assistance in stable isotope and radiocarbon determinations. The authors wish to thank Wally Broecker, Tim Jull and Dietmar Wagenbach, whose constructive and helpful comments improved an earlier version of this paper. This work was partly funded by a Russian foundation for Basic Research (grants 99-05-65075 and 00-05-64736) and by the Program of Integration of Russian Education Ministry (grant 5.1-425).[*AH*]

References

- Yu.K. Vasil'chuk, Oxygen isotope compositions of ground ice application to paleogeocryological reconstructions, vol. 1–2, Theoretical Problems Department, Russian Academy of Sciences and Moscow University Publisher, Moscow, 1992 (in Russian with English comments).
- [2] J. Brown, Radiocarbon dating, Barrow, Alaska, Arctic 18 (1965) 37–48.
- [3] Yu.K. Vasil'chuk, A.C. Vasil'chuk, Oxygen isotope and enzymatic activity variations in the Seyaha syngenetic icewedge complex of the Yamal Peninsula, in: Permafrost, 7th International Conference Proceedings, Yellowknife, Canada, 1998, Collection Nordicana, Centre d'Etudes Nordiques, Université Laval, National Research Council of Canada, pp. 1235–1241.
- [4] Yu.K. Vasil'chuk, A.C. Vasil'chuk, ¹⁴C and ¹⁸O in Siberian syngenetic ice-wedge complexes, Radiocarbon 40 (1998) 883–893.
- [5] J.R. Mackay, Oxygen isotope variations in permafrost, Tuktoyaktuk Peninsula area, Northwest Territories. Geological Survey of Canada, Current Research, Part B, Paper 83-1B, 1983, pp. 67–74.
- [6] W.G. Mook, H.J. Streurman, Physical and chemical aspects of radiocarbon dating, in: W.G. Mook, H.T. Waterbolk (Eds.), Proceedings of the 1st International Sympo-

sium, $^{14}\mathrm{C}$ and Archaology, PACT publications 8, 1983, pp. 31–55.

- [7] S. Wijma, J. van der Plicht, The Groningen AMS tandetron, Nucl. Instrum. Methods Phys. Res. B123 (1997) 218–220.
- [8] K.O. Emery, L.E. Garrison, Sea levels 7000 to 20000 years ago, Science 157 (1967) 684–687.
- [9] T.D. Hamilton, J.L. Craig, P.V. Sellmann, The Fox permafrost tunnel: a Late Quaternary geologic record in central Alaska, Geol. Soc. Am. Bull. 100 (6) (1988) 948– 969.
- [10] A.T. Wilson, ¹⁴C studies of natural ice, Radiocarbon 40 (2) (1998) 953–962.